

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSPTASXS1656

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	MAR 31	IFICDB, IFIPAT, and IFIUDB enhanced with new custom IPC display formats
NEWS	3	MAR 31	CAS REGISTRY enhanced with additional experimental spectra
NEWS	4	MAR 31	CA/CAPplus and CASREACT patent number format for U.S. applications updated
NEWS	5	MAR 31	LPCI now available as a replacement to LDPCI
NEWS	6	MAR 31	EMBASE, EMBAL, and LEMBASE reloaded with enhancements
NEWS	7	APR 04	STN AnaVist, Version 1, to be discontinued
NEWS	8	APR 15	WPIDS, WPINDEX, and WPIX enhanced with new predefined hit display formats
NEWS	9	APR 28	EMBASE Controlled Term thesaurus enhanced
NEWS	10	APR 28	IMSRESEARCH reloaded with enhancements
NEWS	11	MAY 30	INPAFAMDB now available on STN for patent family searching
NEWS	12	MAY 30	DGENE, PCTGEN, and USGENE enhanced with new homology sequence search option
NEWS	13	JUN 06	EPFULL enhanced with 260,000 English abstracts
NEWS	14	JUN 06	KOREAPAT updated with 41,000 documents
NEWS	15	JUN 13	USPATFULL and USPAT2 updated with 11-character patent numbers for U.S. applications
NEWS	16	JUN 19	CAS REGISTRY includes selected substances from web-based collections
NEWS	17	JUN 25	CA/CAPplus and USPAT databases updated with IPC reclassification data
NEWS	18	JUN 30	AEROSPACE enhanced with more than 1 million U.S. patent records
NEWS	19	JUN 30	EMBASE, EMBAL, and LEMBASE updated with additional options to display authors and affiliated organizations
NEWS	20	JUN 30	STN on the Web enhanced with new STN AnaVist Assistant and BLAST plug-in
NEWS	21	JUN 30	STN AnaVist enhanced with database content from EPFULL
NEWS	22	JUL 28	CA/CAPplus patent coverage enhanced
NEWS	23	JUL 28	EPFULL enhanced with additional legal status information from the epoline Register
NEWS	24	JUL 28	IFICDB, IFIPAT, and IFIUDB reloaded with enhancements
NEWS	25	JUL 28	STN Viewer performance improved
NEWS	26	AUG 01	INPADOCDB and INPAFAMDB coverage enhanced

NEWS EXPRESS JUNE 27 08 CURRENT WINDOWS VERSION IS V8.3,
AND CURRENT DISCOVER FILE IS DATED 23 JUNE 2008.

NEWS HOURS STN Operating Hours Plus Help Desk Availability

NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 16:33:28 ON 06 AUG 2008

=> File Medline EMBASE Biosis Caplus
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 16:33:36 ON 06 AUG 2008

FILE 'EMBASE' ENTERED AT 16:33:36 ON 06 AUG 2008
Copyright (c) 2008 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 16:33:36 ON 06 AUG 2008
Copyright (c) 2008 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 16:33:36 ON 06 AUG 2008
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

=> S (raf-1 (3A) RBD) (8A) (variant or mutant or mutated or mutation or mutating or mutagenesis or substitution or substitute or substituted)

L1 2 (RAF-1 (3A) RBD) (8A) (VARIANT OR MUTANT OR MUTATED OR MUTATION OR MUTATING OR MUTAGENESIS OR SUBSTITUTION OR SUBSTITUTE OR SUBSTITUTED)

=> S (gtpas or ras) (8A) (binding affinity)

L2 66 (GTPAS OR RAS) (8A) (BINDING AFFINITY)

=> s l1 and l2

L3 0 L1 AND L2

=> d l1 1-2 bib ab

L1 ANSWER 1 OF 2 EMBASE COPYRIGHT (c) 2008 Elsevier B.V. All rights reserved on STN

AN 1999061407 EMBASE

TI Nuclear magnetic resonance and molecular dynamics studies on the interactions of the Ras-binding domain of Raf-1 with wild-type and mutant Ras proteins.

AU Tateno, Masaru; Ebisuzaki, Toshikazu

CS Computational Science Laboratory, Inst. Phys. and Chem. Res. (RIKEN), 2-1 Hirosawa, Wako-shi, Saitama, 351-0198, Japan.

AU Terada, Tohru; Hashimoto, Kyoko; Yokoyama, Shigeyuki (correspondence)

CS Dept. of Biophysics and Biochemistry, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan.
yokoyama@y-sun.biochem.s.u.-tokyo.ac.jp

AU Smith, Brian O.; Laue, Ernest D.
 CS Cambridge Ctr. for Molec. Recog., Department of Biochemistry, University
 of Cambridge, Tennis Court Road, Cambridge, CB2 1QW, United Kingdom.
 AU Cooper, Jonathan A.
 CS Fred Hutchinson Cancer Res. Center, 1100 Fairview Avenue, North, Seattle,
 WA 98109-1024, United States.
 AU Ito, Yutaka; Shirouzu, Mikako; Kigawa, Takanori; Takio, Koji; Shibata,
 Takehiko
 AU Yokoyama, Shigeyuki (correspondence)
 CS Cellular Signaling Laboratory, Institute Physical Chemical Research, 2-1
 Hirosawa, Wako-shi, Saitama 351-0198, Japan. yokoyama@y-sun.biochem.s.u.-t
 okyo.ac.jp
 SO Journal of Molecular Biology, (12 Feb 1999) Vol. 286, No. 1, pp. 219-232.
 Refs: 74
 ISSN: 0022-2836 CODEN: JMOBAK
 CY United Kingdom
 DT Journal; Article
 FS 029 Clinical and Experimental Biochemistry
 LA English
 SL English
 ED Entered STN: 25 Feb 1999
 Last Updated on STN: 25 Feb 1999
 AB The Ras protein and its homolog, Rap1A, have an identical 'effector
 region' (residues 32-40) preceded by Asp30-Glu31 and Glu30-Lys31,
 respectively. In the complex of the 'Ras-like' E30D/K31E mutant Rap1A
 with the Ras-binding domain (RBD), residues 51-131 of Raf-1, Glu31 in
 Rap1A forms a tight salt bridge with Lys84 in Raf-1. However, we have
 recently found that Raf-1 RBD binding of Ras is indeed reduced by the E31K
 mutation, but is not affected by the E31A mutation. Here, the
 'Rap1A-like' D30E/E31K mutant of Ras was prepared and shown to bind the
 Raf-1 RBD less strongly than wild-type Ras, but slightly more tightly than
 the E31K mutant. The backbone (1)H, (13)C, and (15)N magnetic resonances
 of the Raf-1 RBD were assigned in complexes with the wild-type and
 D30E/E31K mutant Ras proteins in the guanosine 5'-O-(β , γ -
 imidotriphosphate)-bound form. The Lys84 residue in the Raf-1 RBD
 exhibited a large change in chemical shift upon binding wild-type Ras,
 suggesting that Lys84 interacts with wild-type Ras. The D30E/E31K mutant
 of Ras caused nearly the same perturbations in Raf-1 chemical shifts,
 including that of Lys84. We hypothesized that Glu31 in Ras may not be the
 major salt bridge partner of Lys84 in Raf-1. A molecular dynamics
 simulation of a model structure of the Raf-1 RBD.ovrhdot.Ras.ovrhdot.GTP
 complex suggested that Lys84 in Raf-1 might instead form a tight salt
 bridge with Asp33 in Ras. Consistent with this, the D33A mutation
 in Ras greatly reduced its Raf-1 RBD binding
 activity. We conclude that the major salt bridge partner of Lys84 in
 Raf-1 may be Asp33 in Ras.

L1 ANSWER 2 OF 2 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN
 AN 1999:134549 BIOSIS
 DN PREV199900134549
 TI Nuclear magnetic resonance and molecular dynamics studies on the
 interactions of the Ras-binding domain of Raf-1 with wild-type and mutant
 Ras proteins.
 AU Terada, Tohru; Ito, Yutaka; Shirouzu, Mikako; Tateno, Masaru; Hashimoto,
 Kyoko; Kigawa, Takanori; Ebisuzaki, Toshikazu; Takio, Koji; Shibata,
 Takehiko; Yokoyama, Shigeyuki [Reprint author]; Smith, Brian O.; Laue,
 Ernest D.; Cooper, Jonathan A.
 CS Cellular Signaling Lab., Inst. Phys. Chem. Res., 2-1 Hirosawa, Wako-shi,
 Saitama 351-0198, Japan
 SO Journal of Molecular Biology, (Feb. 12, 1999) Vol. 286, No. 1, pp.
 219-232. print.

CODEN: JMOBAK. ISSN: 0022-2836.

DT Article

LA English

ED Entered STN: 31 Mar 1999

Last Updated on STN: 31 Mar 1999

AB The Ras protein and its homolog, Rap1A, have an identical "effector region" (residues 32-40) preceded by Asp30-Glu31 and Glu30-Lys31, respectively. In the complex of the "Ras-like" E30D/K31E mutant Rap1A with the Ras-binding domain (RBD), residues 51-131 of Raf-1, Glu31 in Rap1A forms a tight salt bridge with Lys84 in Raf-1. However, we have recently found that Raf-1 RBD binding of Ras is indeed reduced by the E31K mutation, but is not affected by the E31A mutation. Here, the "Rap1A-like" D30E/E31K mutant of Ras was prepared and shown to bind the Raf-1 RBD less strongly than wild-type Ras, but slightly more tightly than the E31K mutant. The backbone ¹H, ¹³C, and ¹⁵N magnetic resonances of the Raf-1 RBD were assigned in complexes with the wild-type and D30E/E31K mutant Ras proteins in the guanosine 5'-O-(beta,gamma-imidotriphosphate)-bound form. The Lys84 residue in the Raf-1 RBD exhibited a large change in chemical shift upon binding wild-type Ras, suggesting that Lys84 interacts with wild-type Ras. The D30E/E31K mutant of Ras caused nearly the same perturbations in Raf-1 chemical shifts, including that of Lys84. We hypothesized that Glu31 in Ras may not be the major salt bridge partner of Lys84 in Raf-1. A molecular dynamics simulation of a model structure of the Raf-1 RBD:Ras:GTP complex suggested that Lys84 in Raf-1 might instead form a tight salt bridge with Asp33 in Ras. Consistent with this, the D33A mutation in Ras greatly reduced its Raf-1 RBD binding activity. We conclude that the major salt bridge partner of Lys84 in Raf-1 may be Asp33 in Ras.